

Exact Random Walk Distributions using Noncommutative Geometry

Jean Bellissard[†], Carlos J Camacho[‡], Armelle Barelli[†], and Francisco Claro[‡]

[†] Laboratoire de Physique Quantique, UMR5626 associée au CNRS, IRSAMC, Université Paul Sabatier, 118, route de Narbonne, F-31062 Toulouse Cedex 4, France

[‡] Facultad de Física, P. Universidad Católica de Chile, Casilla 306, Santiago 22, Chile

Abstract

Using the results obtained by the non commutative geometry techniques applied to the Harper equation, we derive the areas distribution of random walks of length N on a two-dimensional square lattice for large N , taking into account finite size contributions.

Pacs Numbers : 05.45.+b, 72.15.Qm, 72.10.Bg

Let us consider on a square lattice all closed paths of length N starting at the origin. For such a path Γ , let $A(\Gamma)$ be its algebraic area. As $N \mapsto \infty$, the average size of such a path increases as $\sqrt{N}$ so that $A(\Gamma) \simeq N$ and the renormalized area will be $a = A/N$. We want to compute the probability distribution $\mathcal{P}(A, N)$ of the areas at large but finite N . In the limit $N \mapsto \infty$, the distribution was computed first by Lévy using brownian paths [1]. We will give a method based upon the Harper model, allowing to compute the finite size corrections in a systematic way. The Harper model was designed in 1955 [2] as the simplest non-trivial one describing the motion of an electron sitting on a two-dimensional square lattice and submitted to a uniform magnetic field. Let ϕ be the magnetic flux through the unit cell and let $\phi_0 = h/e$ be the flux quantum. We set $\gamma = 2\pi\phi/\phi_0$. Then given $m = (m_1, m_2) \in \mathbf{Z}^2$, we denote by $W(m)$ the corresponding magnetic translations [3]. They satisfy the Weyl commutation rules

$$W(m)W(m') = W(m + m') \exp(i\frac{\gamma}{2}m \wedge m'), \quad (1)$$

where $m' \wedge m = m'_1 m_2 - m'_2 m_1 \in \mathbf{Z}$. Note that γ plays a rôle similar to the Planck constant in the canonical commutation relation.

Harper's model is given by the following Hamiltonian :

$$H = \sum_{|a|=1} W(a), \quad (2)$$

where $|a| = |a_1| + |a_2|$ if $a = (a_1, a_2) \in \mathbf{Z}^2$. In addition, one defines the trace per unit area as the unique linear map $\mathcal{T}$ on the algebra generated by the $W(m)$'s such that

$$\mathcal{T}(W(m)) = \delta_{m,0} . \quad (3)$$

Then from (1), (2) and (3), we get :

$$\mathcal{T}(H^N) = \sum_{\Gamma : \text{closed paths of length } N} e^{i\gamma A(\Gamma)/2} ,$$

where the sum is taken on the set of closed paths starting at the origin of length N . Note that N should be even to get a non zero sum. Let Ω_N be the number of such closed paths, we then get :

$$\sum_{A=-A_{\max}}^{A_{\max}} \mathcal{P}_N(A/N) \exp(ixA/N) = \Omega_N^{-1} \sum_{\Gamma} \exp(ixA(\Gamma)/N) = \Omega_N^{-1} \mathcal{T}(H^N)|_{\gamma=x/N} . \quad (4)$$

From this relation we obtain :

$$\begin{aligned} \Omega_N = \mathcal{T}(H^N)|_{\gamma=0} &= \int \frac{dk_1 dk_2}{4\pi^2} (2 \cos k_1 + 2 \cos k_2)^N \\ &= \frac{4^{N+1}}{2\pi N} \left(1 + O(1/N^2)\right) \text{ as } N \mapsto \infty . \end{aligned} \quad (5)$$

Moreover as $N \mapsto \infty$, for a given value of x , $\gamma = x/N$ tends to zero, so that we can use a semiclassical argument to compute $\mathcal{T}(H^N)$. It has been shown that the spectrum of H is made of Landau sublevels [4] :

$$E_{\ell}^{\pm}(\gamma) = \pm \left(4 - \gamma(2\ell + 1) + \frac{\gamma^2}{16} [1 + (2\ell + 1)^2] - O(\gamma^3) \right) \quad \ell = 0, 1, \dots, O(1/\gamma) , \quad (6)$$

each with multiplicity per unit area $g_{\ell}^{\pm} = \gamma/2\pi$. So,

$$\mathcal{T}(H^N) = \sum_{\pm} \sum_{\ell} \left(E_{\ell}^{\pm}(\gamma) \right)^N [\gamma/(2\pi)] . \quad (7)$$

This gives :

$$\mathcal{T}(H^N) = \frac{4^{N+1}}{2\pi N} \frac{x/4}{\sinh(x/4)} \left[1 - \frac{1}{2N} \frac{(x/4)^2}{\sinh^2(x/4)} + O(1/N^2) \right] . \quad (8)$$

Using (4), we obtain the probability distribution :

$$\mathcal{P}_N(a) = \frac{\pi}{\cosh^2(2\pi a)} + O(1/N) . \quad (9)$$

We numerically computed $\mathcal{P}_N(a)$ from formula (8) and compared the result with exact numerical calculations (Figure 1).




FIG. 1. Scaling function for the probability of having a loop of area A for a random walk of N steps. Diamond, plus and triangle symbols correspond to the finite size data for $N = 16, 18$ and 20 , respectively. Solid line corresponds to the universal function (9), as $N \mapsto \infty$; whereas dashed lines include the $1/N$ correction term for $N=20$ and 40 .

Acknowledgements : This work has been partially supported by the ECOS/CONY CIT project C94E07 and FONDECYT No. 3940016 (Chile).

References

- [1] Lévy P 1988 *Brownian motion* FW Wiegel
- [2] Harper P G 1955 *Proc. Phys. Soc. Lond. A* **68** 874; *Proc. Phys. Soc. Lond. A* **68** 879
- [3] Zak J 1964 *Phys. Rev. A* **134** 1602 and 1607-1611.
- [4] Rammal R and Bellissard J 1990 *J. Phys. France* **51** 1803